BRAIN RESEARCH MOLECULAR BRAIN RESEARCH

VOL. 50 NOS. 1,2	VO	L.	50	NO	S.	1.2
-------------------------	----	----	----	----	----	-----

CONTENTS

15 OCTOBER 1997

Cited in Biological Abstracts (BIOSIS) – Chem. Abstracts – Index Medicus (MEDLINE) – Current Contents (Life Sci.) – Excerpta Medica (EMBASE) – Psychological Abstracts (PsycINFO) – Pascal et Francis (INIST-CNRS) – RIS (Reference Update) – Current Awareness in Biological Science (CABS)

Research reports

Multiple signaling pathways direct the initiation of tyrosine hydroxylase gene exp	pression in cultured brain neurons	
X. Du and L. Iacovitti (USA)		
Molecular analysis of the monomeric GTP-binding proteins of oligodendrocytes	(200. 0)	
R. Burcelin, A.G. Rodriguez-Gabin, M.J. Charron, G. Almazan and J.N. I		
Activation of CPP-32 protease in hippocampal neurons following ischemia and ep		
F. Gillardon, B. Böttiger, B. Schmitz, M. Zimmermann and KA. Hossmann		1
Controlled targeting of tyrosine hydroxylase protein toward processes of locus co		
L. Bezin, JJ. Diaz, D. Marcel, M. Le Cavorsin, JJ. Madjar, JF. Pujol	and D. Weissmann (France)	2
Identification of cell type-specific promoter elements associated with the rat tyros	ine hydroxylase gene using transgenic founder analysis	
J. Liu, J.P. Merlie, R.D. Todd and K.L. O'Malley (USA)		3
Effects of angiotensinogen antisense oligonucleotides on fluid intake in response	to different dipsogenic stimuli in the rat	
P. Sinnayah, E. Kachab, J. Haralambidis, J.P. Coghlan and M.J. McKinley	(Australia)	4
Evidence for autocrine inhibition of gonadotropin-releasing hormone (GnRH) ger	e transcription by GnRH in hypothalamic GT1-1 neuronal cells	
S. Cho, J. Han, W. Sun, D. Choi, H.B. Kwon, H. Jarry, W. Wuttke and K		5
Selective increase in somatostatin mRNA expression in human basal ganglia in Pa		
D.J. Eve, A.P. Nisbet, A.E. Kingsbury, J. Temlett, C.D. Marsden and O.J.		5
BDNF protection of auditory neurons from cisplatin involves changes in intracellular		
R. Gabaizadeh, H. Staecker, W. Liu and T.R. Van De Water (USA)	7	7
Effects of pentylenetetrazol-induced status epilepticus on c-Fos and HSP72 immur		
J.E. Motte, M.J. Da Silva Fernandes, C. Marescaux and A. Nehlig (France		71
A truncated Reelin protein is produced but not secreted in the 'Orleans' reeler mu		
V. De Bergeyck, K. Nakajima, C. Lambert de Rouvroit, B. Naerhuyzen, A.		
Japan)	Wi. Connect, 1. Milyata, M. Ogawa and K. Mikosinoa (Belgium,	2 /
Adenovirus-mediated expression of AMPA-type glutamate receptor channels in Po		,.
M. Sudo, K. Tsuzuki, H. Okado, A. Miwa and S. Ozawa (Japan)	STZ CERS	
Estrogen regulates galanin but not tyrosine hydroxylase gene expression in the rat		,
		·
J.Y. Tseng, P.E. Kolb, M.A. Raskind and M.A. Miller (USA)	10	/(
Ibuprofen: effect on inducible nitric oxide synthase	10	
N.C. Stratman, D.B. Carter and V.H. Sethy (USA)	10	,
Retinal Muller glia secrete apolipoproteins E and J which are efficiently assemble		
J. Shanmugaratnam, E. Berg, L. Kimerer, R.J. Johnson, A. Amaratunga, B		
Lysophosphatidic acid potentiates ACh receptor currents by G-protein-mediated ac		
T. Nishizaki and K. Sumikawa (Japan, USA)	12	.]
Islet amyloid polypeptide and calcitonin gene-related peptide expression are upreg	ulated in lumbar dorsal root ganglia after unilateral adjuvant-in-	
duced inflammation in the rat paw		
H. Mulder, Y. Zhang, N. Danielsen and F. Sundler (Sweden)	12	7
Regulation of NMDA receptor subunit messenger RNA levels in the rat brain followers		
M.A. Riva, F. Tascedda, E. Lovati and G. Racagni (Italy)	13	6
New isoforms of the chick glutamate receptor subunit GluR4: molecular cloning,		
A. Ravindranathan, T.N. Parks and M.S. Rao (USA)	14	3
Localization of putative calcium-responsive regions in the rat BDNF gene		
J.F. Bishop, G. Joshi, G.P. Mueller and M.M. Mouradian (USA)	15	4
Prostaglandin H synthase-2 and cytosolic phospholipase A ₂ in the hypoxic-ischem		
M. Walton, E. Sirimanne, C. Williams, P.D. Gluckman, J. Keelan, M.D. M.	litchell and M. Dragunow (New Zealand)	5
AP-1 and NF-κB stimulated by carbachol in human neuroblastoma SH-SY5Y cell	s are differentially sensitive to inhibition by lithium	
R.S. Jope and L. Song (USA)	. 17	1
Continuous administration of the glutamate uptake inhibitor L-trans-pyrrolidine-2,4	-dicarboxylate produces striatal lesion	
I-C Liévens M Dutertre C Forni P Salin and I Kerkerian-Le Goff (Fi	rance)	1

(contents continued on p. 322)



0168-328x(19971015)50-1/2-1-7

(contents continued)	
Immunocytochemical identification of pinopsin in pineal glands of chicken and pigeon	
T. Okano, Y. Takanaka, A. Nakamura, K. Hirunagi, A. Adachi, S. Ebihara and Y. Fukada (Japan)	190
Differential expression of sodium channel genes in retinal ganglion cells	
J. Fjell, S. Dib-Hajj, K. Fried, J.A. Black and S.G. Waxman (USA, Sweden)	197
Experimental diffuse axonal injury induces enhanced neuronal C5a receptor mRNA expression in rats	
P.F. Stahel, T. Kossmann, M.C. Morganti-Kossmann, V.H.J. Hans and S.R. Barnum (USA, Switzerland)	205
Different effects of oxidative stress on activation of transcription factors in primary cultured rat neuronal and glial cells	
E. Iwata, M. Asanuma, S. Nishibayashi, Y. Kondo and N. Ogawa (Japan)	213
Dopamine receptor gene expression in an animal model of 'behavioral dependence' on ethanol	
M. Eravci, T. Großpietsch, G. Pinna, O. Schulz, S. Kley, M. Bachmann, J. Wolffgramm, E. Götz, A. Heyne, H. Meinhold and A.	
Baumgartner (Germany)	221
Induction of protein tyrosine phosphatase € transcripts during NGF-induced neuronal differentiation of PC12D cells and during the development of the cerebellum	
Y. Mukouyama, H. Kuroyanagi, T. Shirasawa, T. Tomoda, D. Saffen, M. Oishi and T. Watanabe (Japan)	230
Type 2 interleukin-1 receptor mRNA is induced by kainic acid in the rat brain	
A. Nishiyori, M. Minami, S. Takami and M. Satoh (Japan)	237
Selective effects of partial and severe lesions of the serotonergic systems on Met-enkephalin and substance P neurons in rat basal ganglia	246
V. Compan, P. Salin and A. Daszuta (France)	246
Changes in voltage-dependent calcium channel α_1 -subunit mRNA levels in the kindling model of epileptogenesis	257
H. Hendriksen, W. Kamphuis and F.H. Lopes da Silva (The Netherlands) Activation of heme oxygenase and consequent carbon monoxide formation inhibits the release of arginine vasopressin from rat hypothalamic	231
explants. Molecular linkage between heme catabolism and neuroendocrine function	
C. Mancuso, I. Kostoglou-Athanassiou, M.L. Forsling, A.B. Grossman, P. Preziosi, P. Navarra and G. Minotti (Italy, UK)	267
Molecular characterization of antipeptide antibodies against the 5-HT _{1A} receptor: evidence for state-dependent antibody binding	207
T.E. Anthony and E.C. Azmitia (USA)	277
Regional differences in expression of transcripts for Na ⁺ /Ca ²⁺ exchanger isoforms in rat brain	
L. Yu and R.A. Colvin (USA)	285
Alterations in behavior and opioid gene expression induced by the novel tropane analog WF-31	
J.B. Daunais, S.L. Hart, A. Hedgecock-Rowe, J.J. Matasi, C. Thornley, H.M.L. Davies and L.J. Porrino (USA)	293
mRNA for the m4 muscarinic receptor subtype is expressed in adult rat brain cholinergic neurons	
K. Sugaya, C. Clamp, D. Bryan and M. McKinney (USA)	305
Short communication	
Expression of tyrosine hydroxylase in magnocellular hypothalamic neurons of obese (fa/fa) and lean heterozygous (Fa/fa) Zucker rats	
S. Fetissov, F. Marsais, S. Nicolaïdis and A. Calas (France)	314
Author Index	319